

Sheila¹⁾ reported the influence of the supine position, the sitting position, and the sidelying position for CP of GMFCS V level with an intervention time of 20 min. The results showed that the use of therapeutic positioning in sitting and sidelying positions should be considered as a noninvasive intervention for a population with respiratory compromise.

Taking these results into consideration, even for a short period of intervention time compared with the ARDS case, it is estimated that the redistribution of the expansion gradient from the dorsal side to the ventral side leads to improvement of oxygen supply.

No adverse events were observed in this study during the intervention period. We followed the technique of Langer et al., who manually changed the position using four staff members as a postural change of prone position for ARDS patients. Attention was paid to avoid eye damage and unphysiologic movement of the extremities during the posture changes. In that way, no adverse events related to the prone position were observed during the intervention period¹⁴⁾. Guerin et al. showed all institution that studied the effect of prone position on severe ARDS patients were skilled in the process of turning patients from the supine to the prone position, as shown by the absence of adverse events directly related to repositioning¹²⁾.

Factors considered to have led to the safety of this research were the implementation by 3 to 4 members of the care staff who were familiar with the physical condition of the subjects, consideration for the tracheostomy, monitoring in proximity to the subject, and use of a device for holding the prone position according to the subject's body shape.

In this case, pneumonia and fever were not observed during the intervention period. It was speculated that the natural discharge of respiratory tract secretions in the prone position could have influenced the prevention of aspiration.

In relation to the physical activity and the healthy life span of CP patients, sedentary behavior (SB) is known to have an adverse effect on health¹⁵⁾¹⁶⁾. The metabolic equivalent of a task (MET) used in this table is a physiological measure expressing the energy cost of physical activities and is defined as the ratio of the metabolic rate during a specific physical activity to a reference metabolic rate (1 MET), set by convention at 3.5 mL O₂/kg/min. A recently published standardized definition of SB is any waking behavior characterized by an energy expenditure of 1.5 METs in muscular inactivity while in a sitting or reclining posture¹⁷⁾.

Therefore, strategies to increase physical activity levels in children and adolescents with CP are considered important for long-term health¹⁸⁾. For patients with severe motor disabilities, such as the subjects in this study, who found it difficult to change posture, opportunities for attitude change are considered important exercise opportunities. Therefore, implementation of various posture variations is an important consideration for health life extension.

Although the relationship between postural change and life extension was not clarified because the long-term prognosis was not considered, due to the safety of implementation of prone position therapy, it is desirable to contribute to the prolongation of healthy life in the future.

From the results of this study, prone positioning for quadriplegic patients who present severe motor disabilities due to cerebral disease can be expected to have a certain effect even for short-term intervention, so it is possible to consider safe intervention according to the subjects.

A limitation of this study was that only ten cases were evaluated. Although it was possible to show that effective changes occurred using this method, it is necessary to carry out long-term intervention in the future and verify it including its effects.

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

- 1) Rosenbaum PL, Paneth N, Leviton A, et al: A report: the definition and classification of cerebral palsy. *Dev Med Child Neurol Suppl* 109: 8–14,
- 2) Littleton SR, Heriza CB, Mullenset PA, et al: Effects of positioning on respiratory measures in individuals with cerebral palsy

- and severe scoliosis. *Pediatr Phys Ther* 23: 159—169.
- 3) Reddihough DS, Baikie G, Walstab JE: Cerebral palsy in Victoria, Australia: mortality and causes of death. *J Paediatr Child Health* 37: 183—186.
 - 4) Koulouras V, Papathanakos G, Papathanasiou A, et al: Efficacy of prone position in acute respiratory distress syndrome patients. *World J Crit Care Med* 5: 121—136.
 - 5) Galiatsou E, Kostanti E, Svalna E, et al: Prone position augments recruitment and prevents alveolar overinflation in acute lung injury. *Am J Respir Crit Care Med* 174: 187—197.
 - 6) Mutoh T, Guest RJ, Lamm WJ, et al: Prone position alters the effect of volume overload on regional pleural pressures and improves hypoxemia in pigs in vivo. *Am Rev Respir Dis* 146: 300—306.
 - 7) Palisano RJ, Rosenbaum P, Bartlett D, Livingstone MH: Content validity of the expanded and revised Gross Motor Function Classification System. *Dev Med Child Neurol* 50: 744—750.
 - 8) Bohannon RW, Smith MB: Interrater reliability of a modified Ashworth scale of muscle spasticity. *Phys Ther* 67: 206—207.
 - 9) Ersöz M, Selçuk B, Gündüz R, et al: Decreased chest mobility in children with spastic cerebral palsy. *Turk J Pediatr* 48: 344—350.
 - 10) Park ES, Park JH, Rha DW, et al: Comparison of the ratio of upper to lower chest wall in children with spastic quadriplegic cerebral palsy and normally developed children. *Yonsei Med J* 47: 237—242.
 - 11) Mancebo J, Fernandez R, Blanch L, et al: A multicenter trial of prolonged prone ventilation in severe acute respiratory distress syndrome. *Am J Respir Crit Care Med* 173: 1233—1239.
 - 12) Guerin C, Reignier J, Richard AC, et al: Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med* 368: 2159—2168.
 - 13) Gattinoni L, Tognoni G, Pesenti A, et al: Effect of prone positioning on the survival of patients with acute respiratory failure. *N Engl J Med* 345: 568—573.
 - 14) Langer M, Mascheroni D, Marcolin R, Gattinoni L: The prone position in ARDS patients. A clinical study. *Chest* 94: 103—107.
 - 15) Clanchy KM, Tweedy SM, Boyd R: Measurement of habitual physical activity performance in adolescents with cerebral palsy: a systematic review. *Dev Med Child Neurol* 53: 499—505.
 - 16) Verschuren O, Peterson MD, Leferink S, Darrah J: Muscle activation and energy-requirements for varying postures in children and adolescents with cerebral palsy. *J Pediatr* 165: 1011—1016.
 - 17) Verschuren O, Darrah J, Novak I, et al: Health-Enhancing Physical Activity in Children With Cerebral Palsy: More of the Same Is Not Enough. *Phys Ther* 94: 297—305.
 - 18) Durstine LJ, Painter P, Franklin BA, et al: Physical activity for the chronically ill and disabled. *Sports Med* 30: 207—219.

Reprint request:

Tadashi Matsuda
Department of Rehabilitation, Suita Municipal Disability Support Center I-Hope Suita, 12-27, Senri-Banpaku-Kouen, Suita, Osaka, 565-0826, Japan.

別刷請求先 〒565-0826 大阪府吹田市千里万博公園 12-27
吹田市立障害者支援交流センターあいほうぶ吹
田リハビリテーション部
松田 忠司